

Angen gofal lliniarol heb ei ddiwallu yng Nghymru

Briff polisi

Cynnwys

1. Cyflwyniad	3
2. Beth yw angen gofal lliniarol heb ei ddiwallu?	3
3. Mesur angen gofal lliniarol heb ei ddiwallu	4
4. Lefelau o angen gofal lliniarol heb ei ddiwallu yng Nghymru	4
5. Sut mae angen gofal lliniarol heb ei ddiwallu yn effeithio ar wahanol grwpiau?	5
6. Casgliad	6
7. Atodiad A. Dulliau	7
Ôl-nodiadau	8

Mae'r briff hwn yn edrych ar ymchwil a gynhaliwyd fel rhan o'r *DUECare Project: Defining and Estimating Unmet Palliative Care Needs in the UK**, a arianwyd gan Marie Curie. Arweiniwyd yr ymchwil ar y cyd gan Dr Anna Bone a'r Athro Fliss Murtagh ac mewn partneriaeth gan Sefydliad Cicely Saunders ar gyfer Gofal Lliniarol, Polisi ac Adsefydlu, Coleg y Brenin Llundain, a Chanolfan Ymchwil Gofal Lliniarol Wolfson, Ysgol Feddygol Hull Efrog, Prifysgol Hull, gyda chyfraniadau gan Seicoleg Glinigol, Ysgol Iechyd mewn Gwyddorau Cymdeithasol, Prifysgol Caeredin.

1. Cyflwyniad

MAE gofal lliniarol a diwedd oes yn rhan hanfodol o'n system iechyd a gofal cymdeithasol. Pan fyddwn ni'n ei wneud yn iawn, gall gael effaith ddofn ar bobl sy'n byw gyda salwch terfynol, a'r rhai sy'n agos atynt.

Ond mae gofal diwedd oes ar y dibyn. Mae bylchau yn y ddarpariaeth a phwysau eithriadol ar y system sy'n golygu bod gormod o bobl yn cael eu gadael heb ofal a chymorth angenrheidiol.

Mae'r briff hwn yn tynnu ar ymchwil newydd sydd wedi ceisio mesur yr angen sydd heb ei ddiwallu hwn am ofal lliniarol a gofal diwedd oes. Mae'n nodi faint o bobl yng Nghymru sy'n debygol o fod heb y gofal lliniarol sydd ei angen arnynt nawr, ac yn amcangyfrif sut mae'r sefyllfa hon yn debygol o newid yn y dyfodol.

Yn wyneb y galw cynyddol am ofal lliniarol, allwn ni ddim fforddio gwastraffu amser yn gwneud hyn yn iawn. Mae'r bobl sy'n byw gyda salwch terfynol a'u hanwyliaid yn haeddu gwell.

2. Beth yw angen gofal lliniarol heb ei ddiwallu?

MAE'R ffigurau a gyflwynir yn y briff hwn yn seiliedig ar y diffiniad canlynol o angen heb ei ddiwallu am ofal lliniarol:

Mae anghenion gofal lliniarol heb eu diwallu yn bresennol pan fydd gan berson â salwch sy'n cyfyngu ar fywyd symptomau, pryderon seicogymdeithasol, neu ofynion gofal nad ydynt yn cael sylw digonol drwy'r gwasanaethau sydd ar gael, gydag anallu i gael mynediad at neu dderbyn gofal sy'n canolbwyntio ar y person.

Datblygwyd y diffiniad hwn drwy adolygiad o'r llenyddiaeth a thrafodaethau gweithdy gyda phobl yr effeithir arnynt gan salwch terfynol a'r gweithwyr gofal iechyd a chymdeithasol proffesiynol sy'n gweithio gyda nhw.¹ Mae rhagor o fanylion y dulliau a ddefnyddiwyd i ddatblygu'r diffiniad hwn yn [Atodiad A. Dulliau](#).

3. Mesur angen gofal lliniarol heb ei ddiwallu

ER mwyn asesu nifer a chyfran y bobl sydd ag angen heb ei ddiwallu am ofal lliniarol, rydym wedi defnyddio'r diffiniad uchod o angen heb ei ddiwallu i ddadansoddi canlyniadau arolwg cenedlaethol cynrychioliadol o 1,194 o ofalwyr teuluol sydd wedi colli rhywun, a gomisiynwyd gan Marie Curie ac a gynhaliwyd gan y Swyddfa Ystadegau Gwladol ledled Cymru a Lloegr ym mis Tachwedd 2023.²

Wrth ddadansoddi'r arolwg hwn, fe ystyriom fod gan rywun angen heb ei ddiwallu am ofal lliniarol os oes ganddynt symptomau a phryderon heb eu datrys, **ac** nad ydynt yn derbyn digon o gymorth gan feddygon teulu.

Mae esboniad mwy manwl o'r dulliau a ddefnyddiwyd wedi'i amlinellu yn [Atodiad A. Dulliau](#).

Gan ddefnyddio'r dull uchod, roedd yn bosibl cyfrifo nifer yr ymatebwyr sy'n adrodd bod gan eu hanwyliaid anghenion gofal lliniarol heb eu diwallu yn sampl yr arolwg, ac yn seiliedig ar hyn, amcangyfrif cyfran a nifer y bobl sydd ag angen am ofal lliniarol heb ei ddiwallu yn y boblogaeth ehangach.

Mae hyn wedi sefydlu'r amcangyfrif mwyaf cadarn sydd ar gael o nifer yr oedolion sy'n profi angen heb ei ddiwallu am ofal lliniarol yng Nghymru ar lefel y boblogaeth.

4. Lefelau o angen gofal lliniarol heb ei ddiwallu yng Nghymru

MAE'R canfyddiadau'n dangos bod gan **29% o oedolion yng Nghymru anghenion gofal lliniarol heb eu diwallu**. Mae hynny'n cyfateb i 10,246 o bobl sydd â symptomau a phryderon heb eu datrys, **ac** nad ydynt yn derbyn digon o gymorth gan feddygon teulu ar ddiwedd eu hoes.

Mae'r tabl gyferbyn yn dadansoddi hyn ar draws y ddau fesur sy'n ffurfio ein diffiniad o angen gofal lliniarol heb ei ddiwallu:

Tabl 1: Angen gofal lliniarol heb ei ddiwallu yng Nghymru

Dull o amcangyfrif	Cymru (N=35,511)
1. Symptomau a phryderon	17,209 (49%)
2. Darpariaeth gofal sylfaenol annigonol	15,280 (43%)
3. Amcangyfrif o angen heb ei ddiwallu (1 a 2)	10,246 (29%)

Mae gan Gymru boblogaeth sy'n heneiddio ac o ganlyniad mae'n debygol y bydd yr angen am ofal lliniarol hefyd yn cynyddu yn y blynyddoedd i ddod. Rydym yn amcangyfrif, erbyn y 2040au, y bydd 37,000 o bobl yng Nghymru yn debygol o fod angen gofal lliniarol a gofal diwedd oes bob blwyddyn, cynnydd o 5,000.³

Yn seiliedig ar y dadansoddiad o angen heb ei ddiwallu, amcangyfrifir, heb ymyrraeth ychwanegol, y bydd tua 1,500 yn fwy o bobl yn debygol o wynebu angen gofal lliniarol heb ei ddiwallu yn 2050, o'i gymharu â 2025. Cynnydd o 14%.

5. Sut mae angen gofal lliniarol heb ei ddiwallu yn effeithio ar wahanol grwpiau?

GWYDDYS bod anghydraddoldebau'n bodoli o ran mynediad at wasanaethau iechyd a gofal cymdeithasol, gan gynnwys gofal lliniarol, a phrofiadau ohonynt.

Gan ddefnyddio data ar gyfer Cymru a Lloegr, mae ein dadansoddiad o angen gofal lliniarol heb ei ddiwallu yn dangos amrywiad yn lefelau'r angen heb ei ddiwallu ar sail diagnosis, oedran ac ansicrwydd ariannol. Roedd pobl 85 oed neu'n hŷn yn llai tebygol o fod ag anghenion heb eu diwallu na'r rhai yn yr

ystod oedran 65-84. Mae chwarter (25%) o bobl sy'n byw'n gyfforddus yn profi angen heb ei ddiwallu o'i gymharu â 36% ymhlith y rhai sydd 'prin yn dod i ben neu'n cael pethau'n anodd'. Mae hyn yn golygu bod y rhai sydd 'prin yn dod i ben neu'n cael pethau'n anodd' yn ariannol yn wynebu risg uwch o 45% o angen gofal lliniarol heb ei ddiwallu.⁴

Mae llawer o ffactorau posibl i gyfrif am yr amrywiadau hyn sy'n gofyn am ymchwil pellach i sicrhau bod darparwyr gwasanaethau yn nodi sut i wella gofal.

6. Casgliad

MAE'R ffigurau a gyflwynir yn y briff hwn yn tynnu sylw'n llwyr at y graddau y mae ein system gofal lliniarol a diwedd oes ar y dibyn.

Mae bron i 1 o bob 3 o bobl yng Nghymru heb y gofal a'r gefnogaeth sydd eu hangen arnynt yn ystod misoedd olaf eu bywyd. Mae hyn yn golygu bod miloedd o bobl yn cael eu gadael yn ynysig ac mewn poen.

Gyda'r galw am ofal lliniarol yn debygol o gynyddu a lefelau o angen heb ei ddiwallu yn cael eu rhagweld i dyfu, mae angen gweithredu ar frys i drawsnewid gofal diwedd oes yng Nghymru.

Mae Marie Curie Cymru wedi nodi rhaglen gynhwysfawr o newid polisi yn ein manifesto *Ar fin torri: Amser i drawsnewid gofal diwedd oes yng Nghymru*.⁵

Mae'n darparu cynllun ar gyfer Llywodraeth newydd Cymru yn 5 i sicrhau bod gwasanaethau'n ymatebol i anghenion pobl, bod gofal lliniarol yn cael ei ariannu'n gynaliadwy ac i gryfhau gofal y tu allan i oriau gwaith a gofal cymunedol fel bod gan bobl fynediad 2026 at ofal a chymorth yn eu cartrefi neu'n agos atynt.

Ein her i lunwyr polisi, byrddau iechyd a darparwyr, gan gynnwys ni ein hunain, yw cymryd y camau radical sydd eu hangen i drwsio gofal diwedd oes a sicrhau y gall pawb gael mynediad at y gofal cywir, yn y lle cywir, ar yr amser cywir.

7. Atodiad A. Dulliau

7.1. Diffinio anghenion gofal lliniarol heb ei ddiwallu

Mae'r canfyddiadau ymchwil a gyflwynir yn y briff hwn wedi'u hategu gan waith i ddatblygu diffiniad a rennir o angen heb ei ddiwallu, yn seiliedig ar adolygiad o'r llenyddiaeth a thrafodaethau gweithdy gyda phobl yr effeithir arnynt gan salwch terfynol a'r gweithwyr iechyd a gofal proffesiynol sy'n gweithio gyda nhw.

Nodwyd a graddiwyd nifer o ffactorau a allai gyfrannu at angen heb ei ddiwallu gan gyfranogwyr y gweithdy. Roedd yr elfennau a gafodd y flaenoriaeth fwyaf yn canolbwyntio ar ddarparu gwasanaethau, ansawdd a mynediad, (er enghraifft: diffyg asesiad amserol a chyfannol o symptomau neu ddioddefaint, anallu i gael mynediad at y gwasanaethau sydd eu hangen, a diffyg cydlyniad a pharhad gofal). Blaenoriaethau allweddol eraill a nodwyd oedd diffyg cydnabyddiaeth o anghenion gofal lliniarol, diffyg un pwynt cyswllt ar gyfer cefnogaeth (gan gynnwys gofal y tu allan i oriau), diffyg dilyniant amserol i fynd i'r afael â symptomau a phryderon, gofal medrus a chymwys annigonol, a pheidio â chael eu trin â pharch, urddas ac empathi. Diffyg asesiad amserol a chyfannol o symptomau neu ddioddefaint a gafodd y sgôr cyfanswm uchaf, ac anallu i gael mynediad at y gwasanaethau sydd eu hangen oedd yr eitem a ymddangosodd amlaf yn y 10 uchaf (74%).

Yn seiliedig ar y canfyddiadau hyn, sefydlodd yr ymchwil y diffiniad byr


canlynol ar gyfer angen heb ei ddiwallu am ofal lliniarol:

Mae anghenion gofal lliniarol heb eu diwallu yn bresennol pan fydd gan berson â salwch sy'n cyfyngu ar fywyd symptomau, pryderon seicosymdeithasol, neu ofynion gofal nad ydynt yn cael sylw digonol drwy'r gwasanaethau sydd ar gael, gydag anallu i gael mynediad at neu dderbyn gofal sy'n canolbwyntio ar y person.

7.2. Mesur angen gofal lliniarol heb ei ddiwallu

Mae'r ffigurau yn y briff hwn yn seiliedig ar ddadansoddiad o arolwg cenedlaethol cynrychioliadol o 1,194 o ofalwyr teuluol sydd wedi colli rhywun, a gomisiynwyd gan Marie Curie ac a gynhaliwyd gan y Swyddfa Ystadegau Gwladol ledled Cymru a Lloegr ym mis Tachwedd 2023.

Wrth ddadansoddi'r arolwg hwn, fe ystyriom fod gan rywun angen heb ei ddiwallu am ofal lliniarol os oes ganddynt symptomau a phryderon heb eu datrys, **ac** nad ydynt yn derbyn digon o gymorth gan feddygon teulu ar ddiwedd oes.



Symptomau a phryderon heb eu datrys

Mae'r elfen gyntaf yn seiliedig ar gyfran y bobl â symptomau (corfforol a seicolegol) a phryderon heb eu datrys yn wythnos olaf bywyd gan ddefnyddio'r Raddfa Canlyniadau Gofal Lliniarol Integredig (IPOS). Mae IPOS yn offeryn asesu a ddefnyddir yn rhyngwladol sy'n galluogi cleifion i hunan-adrodd eu profiad o symptomau a phroblemau cyffredin. (Mae fersiwn procsi ar gael pan nad yw cleifion yn gallu ateb cwestiynau eu hunain.)

At ddibenion yr ymchwil hwn, ystyrir bod gan rywun "symptomau a phryderon heb eu trin" os yw cyn-ofalwyr mewn galar wedi nodi bod ganddynt fwy na hanner y sgôr cyfanswm uchaf posibl ar draws pob un

o'r 17 eitem IPOS (≥ 34 allan o 68) yn eu hwythnos olaf o fywyd.

Diffyg mynediad at wasanaethau

Mae'r ail elfen yn seiliedig ar gyfran y bobl sydd heb fynediad at wasanaethau gan ofal sylfaenol i ddatrys symptomau a phryderon.

At ddibenion yr ymchwil hwn, amcangyfrifwyd mynediad annigonol at wasanaethau gofal sylfaenol yn seiliedig ar ymatebion cyn-ofalwyr mewn galar i'r cwestiwn arolwg "Yn y tri mis diwethaf cyn iddynt farw, yn gyffredinol, a ydych chi'n teimlo eu bod wedi cael cymaint o gymorth ag oedd ei angen gan feddygon teulu?"

Ôl-nodiadau

1. A. E. Bone, M. Diggle, T. Johansson, A. Finucane, K. E. Sleeman, J. M. Davies, I. J. Higginson, L. K. Fraser a F.E. Murtagh (2025) "Coproducting a conceptual understanding of unmet palliative care needs: stakeholder workshops using modified nominal group technique" BMC Palliative Care <https://link.springer.com/article/10.1186/s12904-025-01971-4>
2. Cwblhawyd yr arolwg fel rhan o'r prosiect *Better End of Life* (www.mariecurie.org.uk/research-and-policy/policy/better-end-life-report). NB: Mae amrywiad bach yn nifer yr ymatebwyr a adroddwyd rhwng yr adroddiad *Better End of Life* a'r astudiaeth hon. Roedd yr adroddiad *Better End of Life* yn cynnwys data arolwg hyd at fis Medi 2023 gyda chyfanswm o 1,179 o ymatebwyr. Roedd y dadansoddiad a gynhaliwyd ar gyfer prosiect DUECare, y mae'r briff hwn yn seiliedig arno, yn cynnwys data gan ymatebwyr hyd at fis Tachwedd 2023 gyda chyfanswm o 1,194.
3. Dadansoddiad Marie Curie a ddyfynnwyd yn *Ar fin torri: Amser i drawsnewid gofal diwedd oes yng Nghymru*
4. Cyfrifiadau Marie Curie yn seiliedig ar ddata o Brosiect DUECare: *Defining and Estimating Unmet Palliative Care Needs in the UK*
5. mariecurie.org.uk/get-involved/campaigns/senedd-manifesto

* Mae'r briff hwn yn tynnu ar y canlynol: *Defining and measuring unmet palliative care needs* <https://pubmed.ncbi.nlm.nih.gov/41652311/>, *Coproducting a conceptual understanding of unmet palliative care needs* <https://pubmed.ncbi.nlm.nih.gov/41437023/> ac *Unmet palliative care needs in England and Wales: population-based estimates and future projections (2025–2050)*.

Unmet palliative care need in Wales

Policy briefing

February 2026

Contents

1. Introduction	11
2. What is unmet palliative care need?	11
3. Measuring unmet palliative care need	12
4. Levels of unmet palliative care need in Wales	12
5. How are different groups affected by unmet palliative care need?	13
6. Conclusion	14
7. Appendix A. Methods	15
Endnotes	16

The briefing draws on research conducted as part of the *DUECare Project: Defining and Estimating Unmet Palliative Care Needs in the UK**, funded by Marie Curie. The research was co-led by Dr Anna Bone and Professor Fliss Murtagh and carried out in partnership by the Cicely Saunders Institute of Palliative Care, Policy and Rehabilitation, King's College London, and the Wolfson Palliative Care Research Centre, Hull York Medical School, with contributions from the Clinical Psychology, School of Health in Social Science, University of Edinburgh.

1. Introduction

PALLIATIVE and end of life care is a critical part of our health and social care system. When we get it right it can have a profound impact on people living with a terminal illness, and those close to them.

But end of life care is at breaking point. Gaps in provision and a system under severe pressure means too many people are left without the care and support they need.

This briefing draws on new research which has sought to measure this unmet need for palliative and end of life care. It sets out how many people in Wales are likely to be without the palliative care they need now, and estimates how this situation is likely to change in the future.

In the face of growing demand for palliative care we cannot afford to waste time getting this right. People living with a terminal illness and those close to them deserve better.

2. What is unmet palliative care need?

THE figures presented in this briefing are based on the following definition of unmet need for palliative care:

Unmet palliative care needs are present when a person with life-limiting illness has symptoms, psychosocial concerns, or care requirements that are not adequately addressed through available services, with inability to access or receive person-centred care.

This definition was developed through a review of the literature and workshop discussions with people affected by terminal illness and the health and social care professionals who work with them.¹ Further detail on the methods used to develop this definition are outlined in [Appendix A. Methods](#).

3. Measuring unmet palliative care need

In order to assess the number and proportion of people with unmet need for palliative care, we have applied the above definition of unmet need to analyse the results of a nationally representative survey of 1,194 bereaved family carers, commissioned by Marie Curie and conducted by the ONS across England and Wales in November 2023.²

In analysing this survey, we considered someone to have unmet need for palliative care if they have **both** unaddressed symptoms and concerns, **and** do not receive enough help from GPs.

A more detailed explanation of the methods used are outlined in [Appendix A. Methods](#).

Using the method above it was possible to calculate the number of respondents reporting that their loved ones had unmet palliative care needs in the survey sample, and based on this, to estimate the proportion and number of people with unmet need for palliative care in the wider population.

This has established the most robust available estimate of the number of adults who experience unmet need for palliative care in Wales at population level.

4. Levels of unmet palliative care need in Wales

In total the findings indicate that **29% of adults in Wales have unmet palliative care needs**. That is equal to 10,246 people who have both unaddressed symptoms and concerns, and who do not receive enough help from GPs at the end of life.

The table opposite breaks this down across the two measures that form our definition of unmet palliative care need.

Table 1: Unmet palliative care need in Wales

Estimation method	Wales (N=35,511)
1. Symptoms and concerns	17,209 (49%)
2. Insufficient primary care provision	15,280 (43%)
3. Unmet need estimate (1 and 2)	10,246 (29%)

Wales has an ageing population and as a result it is likely that palliative care need will also increase in coming years. We estimate that by the 2040s, 37,000 people in Wales will likely need palliative and end of life care each year, an increase of 5,000.³

Based on the analysis of unmet need, it is estimated that without additional intervention around 1,500 more people are likely to face unmet palliative care need in 2050, compared to 2025. An increase of 14%.

5. How are different groups affected by unmet palliative care need?

INEQUALITIES are known to exist in access to, and experiences of, health and social care services, including palliative care.

Drawing on data for both England and Wales our analysis of unmet palliative care need shows variation in levels of unmet need on the basis of diagnosis, age and financial insecurity. People who were 85 or older were less likely to have unmet needs than those in the 65–84 age range. A quarter (25%) of people living

comfortably experience unmet need compared to 36% among those ‘just about getting by or finding things difficult’. This means those ‘just about getting by or finding things difficult’ financially faced an increased risk of unmet palliative care need of 45%.⁴

There are many potential factors for these variations that require further research to ensure service providers identify how to improve care.

6. Conclusion

THE figures presented in this briefing bring into stark focus the extent to which our palliative and end of life care system is at breaking point.

Almost 1 in 3 people in Wales are without the care and support they need in their final months of life. This means thousands of people are being left isolated and in pain.

With the demand for palliative care likely to increase and levels of unmet need predicted to grow, urgent action is needed to transform end of life care in Wales.

Marie Curie Cymru has set out a comprehensive programme of policy change in our manifesto *At breaking point: Time to transform end of life care in Wales*.⁵

It provides a roadmap for the incoming Welsh Government in 2026 to ensure that services are responsive to people's needs, that palliative care is sustainably funded and to strengthen out of hours and community care so people have 24/7 access to care and support at or close to home.

Our challenge to policymakers, health boards and providers, including ourselves, is to take the radical action needed to fix end of life care and ensure that everyone can access the right care, in the right place, at the right time.

7. Appendix A. Methods

7.1. Defining unmet palliative care need

The research findings presented in this briefing are underpinned by work to develop a shared definition of unmet need, based on a review of the literature and workshop discussions with people affected by terminal illness and the health and care professionals who work with them.

A number of factors that may contribute to unmet need were identified and ranked by workshop participants. The most highly prioritised elements focused on service delivery, quality and access, (for example: lack of timely and holistic assessment of symptoms or suffering, inability to access services needed, and lack of coordination and continuity of care). Other key priorities identified were the lack of recognition of palliative care needs, absence of a single point of contact for support (including out-of-hours care), lack of timely follow up to address symptoms and concerns, insufficient skilled and competent care, and not being treated with respect, dignity, and empathy. Lack of timely and holistic assessment of symptoms or suffering received the highest total score, and inability to access services needed was the item most frequently appearing in the top 10 (74%).


Based on these findings, the research established the following short definition for unmet need for palliative care:

Unmet palliative care needs are present when a person with life-limiting illness has symptoms, psychosocial concerns, or care requirements that are not adequately addressed through available services, with inability to access or receive person-centred care.

7.2. Measuring unmet palliative care need

The figures in this briefing are based on analysis of a nationally representative survey of 1,194 bereaved family carers, commissioned by Marie Curie and conducted by the ONS across England and Wales in November 2023.

In analysing this survey we considered someone to have unmet need for palliative care if they have **both** unaddressed symptoms and concerns, **and** do not receive enough help from GPs at the end of life.



Unaddressed symptoms and concerns

The first element is based on the proportion of people with unaddressed symptoms (physical and psychological) and concerns in the last week of life using the Integrated Palliative Care Outcome Scale (IPOS). IPOS is an internationally used assessment tool that enables patients to self-report their experience of common symptoms and issues. (A proxy version is available when patients are unable to answer questions themselves).

For the purposes of this research someone is considered to have “unaddressed symptoms and concerns” if bereaved former carers reported that they had more

than half the maximum possible total score across all 17 IPOS items (≥ 34 out of 68) in their final week of life.

Lack of access to services

The second element is based on the proportion of people lacking access to services from primary care to resolve symptoms and concerns.

For the purposes of this research inadequate access to basic primary care services was estimated based on bereaved former carers’ responses to the survey question “In the last three months before they died, overall, do you feel they got as much help as needed from GPs?”

Endnotes

1. A. E. Bone, M. Diggle, T. Johansson, A. Finucane, K. E. Sleeman, J. M. Davies, I. J. Higginson, L. K. Fraser and F. E. Murtagh (2025) “Coproducing a conceptual understanding of unmet palliative care needs: stakeholder workshops using modified nominal group technique” BMC Palliative Care <https://link.springer.com/article/10.1186/s12904-025-01971-4>
2. Survey completed as part of the Better End of Life project (mariecurie.org.uk/research-and-policy/policy/better-end-of-life-report). NB: There is a small variation in the reported number of respondents between the *Better End of Life* report and this study. The *Better End of Life* report included survey data to September 2023
3. Marie Curie analysis cited in *At breaking point: Time to transform end of life care in Wales*
4. Marie Curie calculations based on data from the *DUECare Project: Defining and Estimating Unmet Palliative Care Needs in the UK*
5. mariecurie.org.uk/get-involved/campaigns/senedd-manifesto

* This briefing draws on the following: *Defining and measuring unmet palliative care needs* <https://pubmed.ncbi.nlm.nih.gov/41652311/>, *Coproducing a conceptual understanding of unmet palliative care needs* <https://pubmed.ncbi.nlm.nih.gov/41437023/> and *Unmet palliative care needs in England and Wales: population-based estimates and future projections (2025-2050)*